

This paper is dedicated  
to the memory of Juergen Moser.

P.G.Grinevich<sup>1</sup> S.P.Novikov<sup>2</sup>

# Topological Charge of the real periodic finite-gap Sine-Gordon solutions

## Contents

|          |                                                                                                                     |           |
|----------|---------------------------------------------------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction. Reality Problems for the Algebro-Geometric Solutions.</b>                                          | <b>2</b>  |
| <b>2</b> | <b>Finite-gap (algebro-geometrical) solutions to the Sine-Gordon equation. Known results.</b>                       | <b>3</b>  |
| 2.1      | The Sine-Gordon equation. Zero Curvature representation. . .                                                        | 3         |
| 2.2      | Algebro-geometric solutions: The complex theory. . . . .                                                            | 6         |
| 2.3      | Real finite-gap SG solutions: known results. . . . .                                                                | 8         |
| <b>3</b> | <b>Effective description of the real tori. Admissible polynomials</b>                                               | <b>10</b> |
| <b>4</b> | <b>Topological charge.</b>                                                                                          | <b>15</b> |
| <b>5</b> | <b>Appendix 1. The Cherednik differential for the generic periodic real SG solutions.</b>                           | <b>21</b> |
| <b>6</b> | <b>Appendix 2. Numerical simulations.</b>                                                                           | <b>23</b> |
| <b>7</b> | <b>Appendix 3. Fourier Transform on the Riemann Surfaces: The continuous analog of the Krichever-Novikov bases.</b> | <b>24</b> |

---

<sup>1</sup>Landau Institute for Theoretical Physics of the Russian Academy of Sciences, Moscow 117940 Kosygin Street 2, e-mail pgg@landau.ac.ru. This work is partially supported by the RFBR Grant No 98-01-01161 and INTAS Grant No 99-01782. Numerical simulations were partially performed on a computer, donated to the author by the Humboldt Foundation.

<sup>2</sup>IPST, University of Maryland-College Park, MD 20742-2431 and Landau Institute for Theoretical Physics, Moscow, e-mail novikov@ipst.umd.edu. This work is partially supported by the NSF Grant DMS-0072700.

# 1 Introduction. Reality Problems for the Algebro-Geometric Solutions.

The so-called finite-gap or algebro-geometric solutions were found for KdV in the work [1] in 1974. Their complete theory involving the spectral theory of periodic (quasiperiodic) finite-gap Schrödinger operators on the line and its unification with algebraic geometry and analysis on the Riemann surfaces, time dynamics and algebro-geometric hamiltonian aspects was constructed in the works [2, 3, 4, 5, 6, 7] (see in the survey article [8]) and in the works [9, 10].

It was extended in many works to a huge number of 1+1 systems. In the work [11] a purely algebraic procedure cleaned from the spectral theory was formulated on the basis of these works and applied to the 2 + 1 systems like KP. The algebro-geometric spectral theory of the stationary periodic 2D Schrödinger operator was developed in the works [12, 13]. Some classical higher rank problems for the commuting OD operators were solved (see [14]).

However, the algebraic methods free from the spectral theory have a weak point in applications: it is very difficult in many cases to select effectively the proper “real” (i.e. physically or geometrically meaningful) solutions among the complex solutions given by the  $\Theta$ -functional expressions. Some physical properties may be completely invisible from these formulas. One may observe that the effective solution of this problem is normally easy exactly if corresponding Lax operator  $L$  (or its  $\lambda$ -dependent analog) is self-adjoint or has a good spectral theory in the Hilbert space anyway, and difficult otherwise.

The reality problem is trivial for KdV. It is also trivial for the defocusing Nonlinear Schrödinger equation  $NLS_+$  and for Sinh-Gordon equation

$$\frac{\partial^2 u}{\partial \xi \partial \eta} = 4 \sinh u(\xi, \eta).$$

However, the reality problem is nontrivial for such famous classical systems as the focusing Nonlinear Schrödinger  $NLS_-$  and the Sine-Gordon (SG) system.

$$u_{tt} - u_{xx} + \sin u(x, t) = 0$$

For example, until now there was no formula calculating such elementary physical characteristic of the real finite-gap (algebro-geometric) solutions periodic in the space variable  $x$  with period  $T$  as the so-called **Topological Charge**. Let us remind that we call function  $u(x)$  periodic in this case if

$\exp\{iu(x+T)\} = \exp\{iu(x)\}$ . The topological charge of this function is equal to the integer number  $n = [u(x+T) - u(x)]/2\pi$ . The density of the topological charge is equal to the real number  $n/T$ . It can be extended to all quasiperiodic functions in the variable  $x$ . This quantity survives all real periodic nonintegrable perturbations. Therefore it is especially valuable in applications.

Nobody was able to find out what the description of the real components described in terms of the Jacoby torus and  $\Theta$ -functions (found in the works [15, 16]) implies for the values of the topological charge.

**Problem:** How to calculate the topological charge of real periodic SG solutions in terms of the Spectral Data: the Riemann surfaces and divisors.

An attempt has been made almost 20 years ago in the work [17] to develop a method for the solution of this problem. This method has been called an “Algebro-Topological” approach to the reality problems. An interesting formula has been proposed. However, as it was pointed out in [18], the idea of the proof presented in [17] failed. In this work we found a new development of the main idea of the work [17]. We present here a full proof of the formula for the topological charge based on the new effective description of the real components in terms of divisors. A partial case of this result is announced by the authors in the short note [19] where some ideas are briefly described.

In the next work we are going to publish also a correction and a full proof of the formula for the so-called Action Variables announced in the work [17] (the standard field-theoretical Poisson Bracket restricted on the space of finite-gap real SG solutions is the most interesting nontrivial case here).

## 2 Finite-gap (algebro-geometrical) solutions to the Sine-Gordon equation. Known results.

### 2.1 The Sine-Gordon equation. Zero Curvature representation.

The famous Sine-Gordon equation (SG)

$$u_{tt} - u_{xx} + \sin u(x, t) = 0 \tag{1}$$

appeared in the XIX century Geometry in the light-cone representation

$$u_{\xi\eta} = 4 \sin u, \quad u = u(\xi, \eta), \quad (2)$$

where

$$x = 2(\xi + \eta), \quad t = 2(\xi - \eta), \quad \partial_\xi = 2\partial_x + 2\partial_t, \quad \partial_\eta = 2\partial_x - 2\partial_t. \quad (3)$$

A broad family of the “soliton-type solutions” was found already in XIX century: it was based on the discovery by Bianchi and S.Lie the substitution called later the “Backlund Transformation”.

The analogy of SG with KdV (i.e. association with the so-called “Inverse Scattering Transform”) was found in the early 1970s by G.Lamb (see [20]). The modern approach to the study SG equation was developed in the work [21] in 1974. It is based on the representation of the SG equation as a consistency condition for the following linear systems rationally dependent on the parameter  $\lambda$ :

$$\Psi_\xi = U\Psi, \quad \Psi_\eta = V\Psi, \quad (4)$$

where

$$\Psi = \Psi(\lambda, \xi, \eta) = \begin{pmatrix} \psi_1(\lambda, \xi, \eta) \\ \psi_2(\lambda, \xi, \eta) \end{pmatrix}, \quad (5)$$

$$U = U(\lambda, \xi, \eta) = \begin{bmatrix} \frac{i u_\xi}{2} & 1 \\ -\lambda & -\frac{i u_\xi}{2} \end{bmatrix}, \quad V = V(\lambda, \xi, \eta) = \begin{bmatrix} 0 & -\frac{1}{\lambda} e^{iu} \\ e^{-iu} & 0 \end{bmatrix} \quad (6)$$

In the variables  $(x, t)$  we have:

$$\Psi_x = \frac{1}{4}(U + V)\Psi, \quad \Psi_t = \frac{1}{4}(U - V)\Psi, \quad (7)$$

where

$$U = U(\lambda, x, t) = \begin{bmatrix} i(u_x + u_t) & 1 \\ -\lambda & -i(u_x + u_t) \end{bmatrix}, \quad (8)$$

$$V = V(\lambda, x, t) = \begin{bmatrix} 0 & -\frac{1}{\lambda} e^{iu} \\ e^{-iu} & 0 \end{bmatrix}. \quad (9)$$

**Definition 1** We call the compatibility condition for pair of linear systems like above (see (4), (7)) a (rationally)  $\lambda$ -**dependent Zero Curvature representation**. It can be written in algebraic form

$$[\partial_\xi - U(\lambda), \partial_\eta - V(\lambda)] = 0.$$

Let us mention that the standard Lax representation for the KdV and all higher KdV equations  $dL/dt_n = [A_n, L]$  based on the  $\lambda$ -independent linear differential operators  $L, A_n$ , has been transformed into the (polynomially)  $\lambda$ -dependent zero-curvature representation in the work [1] for the needs of the periodic problem (to study the finite-gap solutions invented in this work). Very interesting elliptic zero curvature representations appeared later. Until now we do not know interesting effectively written zero-curvature representations with more complicated  $\lambda$ -dependence.

**Definition 2** A SG solution  $u(x, t)$  is called **periodic** in  $x$  if the quantity  $e^{iu}$  is periodic with period  $T$  in the real variable  $x \in \mathbb{R}$ .

**Definition 3** The quantity  $n = [u(x + T) - u(x)]/2\pi$ , where  $T$  is the period is called **topological charge**. The ratio  $\bar{n} = n/T$  is called **density of topological charge**.

The following statement is well-known in the theory of functions:

**Lemma 1** Let  $u(\vec{X})$ ,  $X \in \mathbb{R}^n$  be a smooth function in  $\mathbb{R}^n$  such, that  $\exp(iu(\vec{X}))$  is single-valued on the torus  $\mathbb{R}^n/\mathbb{Z}^n$ , i.e.  $\exp(iu(\vec{X} + \vec{N})) = \exp(iu(\vec{X}))$  for any integer vector  $\vec{N}$ . Denote by  $u(x)$  restriction of  $u(\vec{X})$  to the strait line  $\vec{X} = \vec{X}_0 + x \cdot \vec{v}$ . Then the density of topological charge  $\bar{n} = \lim_{T \rightarrow \infty} [u(x + T) - u(x)]/2\pi T$  is well-defined; it does not depend on the point  $\vec{X}_0$  and can be expressed by the following formula:

$$\bar{n} = \sum_{k=1}^n n_k v^k, \quad (10)$$

where  $\vec{v} = (v^1, v^2, \dots, v^n)$ , and  $n_k$  are topological charges along the basic cycles  $\mathcal{A}_k$ ,  $k = 1, \dots, n$ :

$$u(X^1, X^2, \dots, X^k + 1, \dots, X^n) - u(X^1, X^2, \dots, X^k, \dots, X^n) = 2\pi n_k. \quad (11)$$

The **Topological Charge along the cycle**  $\mathcal{A} = \sum l_k \mathcal{A}_k \in H_1(T^n, \mathbb{Z})$  is equal to the linear combination  $n(\mathcal{A}) = \sum l_k n_k$  by definition.

## 2.2 Algebro-geometric solutions: The complex theory.

Let us recall the construction of general complex “finite-gap” SG solutions. They were constructed almost immediately after the finite-gap KdV theory (see [23, 24]).

Take the following “spectral data” similar to KdV:

1. A nonsingular hyperelliptic Riemann surface  $\Gamma$  [ $\mu^2 = R(\lambda)$ ], where  $R(\lambda) = \prod_{k=0}^{2g} (\lambda - E_k)$ , such that  $E_0 = 0$  and  $E_i \neq E_j, i, j = 0, 1, \dots, 2g$ . It has exactly  $2g + 2$  branching points  $E_0 = 0, E_1, \dots, E_{2g}, \infty$ ; the genus of  $\Gamma$  is equal to  $g$ . A point  $\gamma \in \Gamma$  is by definition a pair of complex numbers  $\gamma = (\lambda, \mu)$  such, that  $\mu^2 = R(\lambda)$ .
2. A divisor  $D$  of degree  $g$ , i.e. set (or formal sum) of  $g$  points  $D = \gamma_1 + \dots + \gamma_g$ .

For generic data  $\Gamma, D$  there exists an unique two-component “Baker-Akhiezer” vector-function  $\Psi(\gamma, \xi, \eta)$  such that

1. For fixed  $(\xi, \eta)$  the function  $\Psi(\gamma, \xi, \eta)$  is meromorphic in the variable  $\gamma \in \Gamma$  outside the points  $0, \infty$  and has at most 1-st order poles at the divisor points  $\gamma_k, k = 1, \dots, g$ .
2.  $\Psi(\gamma, \xi, \eta)$  has essential singularities at the points  $0, \infty$  with the following asymptotics:

$$\Psi(\gamma, \xi, \eta) = \begin{pmatrix} 1 + o(1) \\ i\sqrt{\lambda} + O(1) \end{pmatrix} e^{i\sqrt{\lambda}\xi} \quad \text{as } \lambda \rightarrow \infty, \quad (12)$$

$$\Psi(\gamma, \xi, \eta) = \begin{pmatrix} \phi_1(\xi, \eta) + o(1) \\ i\sqrt{\lambda}\phi_2(\xi, \eta) + O(\lambda) \end{pmatrix} e^{-\frac{i}{\sqrt{\lambda}}\eta} \quad \text{as } \lambda \rightarrow 0, \quad (13)$$

where  $\phi_1(\xi, \eta), \phi_2(\xi, \eta)$  are some functions of the variables  $\xi, \eta$ .

Let us point out that the divisor of zeroes of the first component  $\psi_1$  we denote  $D(\xi, \eta) = \sum_j \gamma_j(\xi, \eta)$ . For  $\xi = 0, \eta = 0$  we have  $D(0, 0) = D$ . The equations for this divisor in the variables  $\xi, \eta$  we call Dubrovin equations. They are important for our work (see below). The vector-function  $\Psi(\gamma, \xi, \eta)$  satisfies to the zero-curvature equations (4) with potential given by the formula

$$u(\xi, \eta) = i \ln \frac{\phi_2(\xi, \eta)}{\phi_1(\xi, \eta)}. \quad (14)$$

Therefore the function  $u(\xi, \eta)$  solves the SG equation (2).

Both functions  $\Psi$  and  $u$  can be expressed through the  $\Theta$ -functions associated with Riemann surface  $\Gamma$ . It is easier to express the solution  $u$  through the  $\Theta$ -functions than  $\Psi$ . Let us mention that the first  $\Theta$ -functional formula for the functions like  $\Psi$  was obtained by A. Its in the case of KdV; it was found later than the Its-Matveev formula for the potential and KdV solution  $u(x, t)$ —see the [8], Appendix 1. In the higher rank problems there exist sometimes explicit formulas for the functions like  $u$ , but  $\Psi$  cannot be found explicitly (see [14]). As a second remark let us point out that the first component  $\psi_1$  of the Baker-Akhiezer vector-function  $\Psi$  is a partial case of the general scalar two-point Baker-Akhiezer functions satisfying to the second order linear Schrödinger equation  $L_1\psi_1 = 0$  invented in the work [12]. Here we have

$$L_1 = \frac{\partial^2}{\partial \xi \partial \eta} + A(\xi, \eta) \frac{\partial}{\partial \eta} + W(\xi, \eta),$$

$-A = \partial_\xi \log \phi_1$ . In this special case our surface  $\Gamma$  is hyperelliptic, the selected points coincide with  $0, \infty$  and corresponding local parameters are chosen as  $\sqrt{\lambda}, 1/\sqrt{\lambda}$ . The second function  $\psi_2$  satisfies to similar equation  $L_2\psi_2 = 0$  where the operator  $L_2$  is obtained from  $L_1$  by the so-called Laplace Transformations and vice versa. It is known that cyclic Laplace Chains of the length 2 lead to the SG equation in the general complex case (see in the work [22]).

For the generic “spectral data”  $\Gamma$ ,  $D$  corresponding SG solutions are complex and may have singularities for real  $x$ . The function  $\exp\{iu\}$  can be naturally considered as a certain meromorphic function on the complex Jacoby torus  $T^{2g} = J(\Gamma)$  restricted to a linear subspace.

For the effective reconstruction of the function  $u(\xi, \eta)$  we use the formula

$$e^{iu(\xi, \eta)} = \frac{\prod_{j=0}^g (-\lambda_j(\xi, \eta))}{\sqrt{\prod_{j=1}^{2g} E_j}} \quad (15)$$

Dubrovin equations in the variables  $\xi, \eta$  for the divisor  $D(\xi, \eta)$  of zeroes of the first function  $\psi_1$ , where  $D(\xi, \eta) = \gamma_1(\xi, \eta) + \dots + \gamma_g(\xi, \eta)$  and  $\gamma_j(\xi, \eta) = (\lambda_j(\xi, \eta), \mu_j(\xi, \eta))$ , have the form:

$$\left\{ \begin{array}{l} \frac{\partial \lambda_k(\xi, \eta)}{\partial \xi} = -2i \frac{\mu_k(\xi, \eta)}{\prod_{j \neq k} (\lambda_k(\xi, \eta) - \lambda_j(\xi, \eta))} \\ \frac{\partial \mu_k(\xi, \eta)}{\partial \xi} = -i \frac{R'(\lambda_k(\xi, \eta))}{\prod_{j \neq k} (\lambda_k(\xi, \eta) - \lambda_j(\xi, \eta))} \\ \frac{\partial u(\xi, \eta)}{\partial \xi} = -2 \sum_k \frac{\mu_k(\xi, \eta)}{\lambda_k(\xi, \eta) \prod_{j \neq k} (\lambda_k(\xi, \eta) - \lambda_j(\xi, \eta))}, \end{array} \right. \quad (16)$$

where

$$R'(\lambda) = \frac{\partial R(\lambda)}{\partial \lambda} \quad (17)$$

$$\left\{ \begin{array}{l} \frac{\partial \lambda_k(\xi, \eta)}{\partial \eta} = 2i \frac{\mu_k(\xi, \eta) e^{iu(\xi, \eta)}}{\lambda_k(\xi, \eta) \prod_{j \neq k} (\lambda_k(\xi, \eta) - \lambda_j(\xi, \eta))} \\ \frac{\partial \mu_k(\xi, \eta)}{\partial \eta} = i \frac{R'(\lambda_k(\xi, \eta)) e^{iu(\xi, \eta)}}{\lambda_k(\xi, \eta) \prod_{j \neq k} (\lambda_k(\xi, \eta) - \lambda_j(\xi, \eta))} \\ \frac{\partial u(\xi, \eta)}{\partial \eta} = 2 \sum_k \frac{\mu_k(\xi, \eta) e^{iu(\xi, \eta)}}{\lambda_k(\xi, \eta)^2 \prod_{j \neq k} (\lambda_k(\xi, \eta) - \lambda_j(\xi, \eta))}, \end{array} \right. \quad (18)$$

$e^{iu}$  is defined by (15).

### 2.3 Real finite-gap SG solutions: known results.

It turns out that the simplest problem here is to find the class of Riemann surfaces corresponding to the real solutions. This class was found in the work [24]:

The set of branching points of the Riemann surface  $\Gamma$  should contain real points or complex conjugate pairs of points only. All real branching points should be nonpositive. Therefore we may assume, that the first  $2m + 1$  of



them  $E_0 = 0$ ,  $E_1, E_2, \dots, E_{2m}$  are real and  $0 > E_1 > E_2 > \dots > E_{2m}$ . For  $k > m$  we have  $E_{2k} = \overline{E_{2k-1}}$ ,  $\text{Im } E_{2k} \neq 0$ . Therefore our Riemann surface (algebraic curve)  $\Gamma$  is real and admits a  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -group of involutions generated by the standard holomorphic involution  $\sigma : \Gamma \rightarrow \Gamma$  such that

$$\sigma(\lambda, \mu) = (\lambda, -\mu), \quad (19)$$

and by the antiholomorphic involution  $\tau$  such that  $\tau_1 = \sigma\tau = \tau\sigma$ :

$$\tau(\lambda, \mu) = (\bar{\lambda}, \bar{\mu}). \quad (20)$$

Which divisors on these surfaces generate real SG solutions?

They were found in the work [25]. Let us formulate this result.

Take a real Riemann surface of the class described above. Try to construct a meromorphic differential  $\Omega$  with two simple poles located in the points  $0, \infty$  and with  $2g$  zeroes located in the points  $D + \tau D$  where  $D = \gamma_1 + \dots, \gamma_g$ .

**Definition 4 .** *We call the divisor  $D$  admissible if and only if such differential  $\Omega$  exists.*

The admissible divisors lead to the real nonsingular solutions of the SG equation. According to the result of [25] any real solution is nonsingular. So the property of divisor to be admissible is sufficient for the solution to be real and smooth quasiperiodic function for the nonsingular real algebraic curve  $\Gamma$ .

We shall explain below (see Appendix 1) that this requirement is also necessary for the periodic solution to be real nonsingular.

Let us demonstrate here a simple proof of the statement:

The potential  $u(\xi, \eta)$  is real if the condition formulated above is satisfied

Proof. Consider the following 1-forms:

$$\Omega_1 = \psi_1(\gamma, \xi, \eta) \overline{\psi_1(\tau\gamma, \xi, \eta)} \Omega, \quad \Omega_2 = \frac{-1}{\lambda} \psi_2(\gamma, \xi, \eta) \overline{\psi_2(\tau\gamma, \xi, \eta)} \Omega. \quad (21)$$

We may assume, that the residues of the form  $\Omega$  are equal to the number  $\pm 1$  at the points  $0, \infty$  respectively. Then the residues of the differentials  $\Omega_1, \Omega_2$  at the point  $\infty$  are equal to the number  $-1$ . Calculating the residues at the point  $0$ , we get:

$$1 = \phi_1(\xi, \eta) \overline{\phi_1(\xi, \eta)} = \phi_2(\xi, \eta) \overline{\phi_2(\xi, \eta)}. \quad (22)$$

Therefore we have

$$\left| \frac{\phi_2(\xi, \eta)}{\phi_1(\xi, \eta)} \right| = 1 \quad (23)$$

and the potential  $u(\xi, \eta)$  in (14) is real. Our statement is proved.

The form  $\Omega_1$  defined above has simple poles at the points  $0, \infty$  and zeroes at the points  $\gamma_1(\xi, \eta), \dots, \gamma_g(\xi, \eta), \tau\gamma_1(\xi, \eta), \dots, \tau\gamma_g(\xi, \eta)$ . Therefore the divisor of zeroes of  $\psi_1$   $D(\xi, \eta) = \gamma_1(\xi, \eta) + \dots + \gamma_g(\xi, \eta)$  is admissible for all  $\xi, \eta$ .

In the work [25] the number of real components was found: it is equal to  $2^m$  where  $2m$  is a number of the strictly negative branching points. No effective description of these components was presented in this work. In the later works [15],[16] all real components were found and described in terms of Jacobian variety associated with  $\Theta$ -functions. It became clear that this description is very hard to use for the studying simple fundamental properties of solutions. For example, this description do not imply any formula for the topological charge. A most recent survey of this subject can be found in [26].

In the joint work [17] an attempt has been made to describe real components in homological terms involving the cycles  $[\gamma_j] \in H_1(\Gamma \setminus (0 \cup \infty), Z)$  covered by the moving zeroes  $\gamma_j(x)$  of the Baker-Akhiezer function  $\psi_1$  on the Riemann surface  $\Gamma$ . In the special parts of the space of parameters this idea led to the interesting formulas for the topological charge and action variables in [17]. However, an attempt to extend it to the whole phase space failed as it mentioned in [18]. In the present work we are going to prove formula for the topological charge proposed in [17, 18].

### 3 Effective description of the real tori. Admissible polynomials

Let us consider only real Riemann surfaces of the type already described above. To construct a real solution to the SG equation we need to choose a meromorphic differential  $\Omega$  described in the previous paragraph.

**Lemma 2** *The differential  $\Omega$  can be expressed in the following form*

$$\Omega = \left( 1 - \frac{\lambda P_{g-1}(\lambda)}{R(\lambda)^{1/2}} \right) \frac{d\lambda}{2\lambda}, \quad (24)$$

Here  $P_{g-1}(\lambda)$  is a polynomial of degree at most  $g - 1$  with real coefficients such, that the system

$$\begin{cases} \mu = \lambda P_{g-1}(\lambda) \\ R(\lambda) = \mu^2, \end{cases} \quad (25)$$

has solutions exactly in the points  $(0, 0), \gamma_1, \dots, \gamma_g, \tau\gamma_1, \dots, \tau\gamma_g$ .

As it was already mentioned, the proof of this lemma immediately follows from the general classification of the meromorphic differentials on the surface  $\Gamma$ . We simply represented it in this standard form.

**Definition 5 .** *We call a real polynomial  $P_{g-1}$  of the degree at most  $g - 1$  admissible if there exists a divisor  $D = \gamma_1 + \dots + \gamma_g$  such the system (25) has solutions in the points  $(0, 0)$  and  $D + \tau D$ . In particular, any real solution of this system (25) except  $(0, 0)$  should have even multiplicity.*

Admissible polynomials  $P_{g-1}(\lambda)$  can be characterized in the following way:

Consider a pair of functions of the real variable  $\lambda$ :

$$f_{\pm}(\lambda) = \pm \frac{\sqrt{R(\lambda)}}{\lambda}. \quad (26)$$

We assume, that the functions  $f_{\pm}(\lambda)$  are defined only for such  $\lambda \neq 0$  that  $R(\lambda) \geq 0$ , i.e. these functions are real-valued. Let us draw the graphs of these functions  $y = f_+(\lambda)$ ,  $y = f_-(\lambda)$ , and fill in the following domains by the black color:

$$\begin{aligned} \lambda < 0, y^2 &< \frac{R(\lambda)}{\lambda^2}, \\ \lambda > 0, y^2 &> \frac{R(\lambda)}{\lambda^2}. \end{aligned} \quad (27)$$




Fig 1.

**Lemma 3** *The polynomial  $P_{g-1}(\lambda)$  is admissible if and only if the graph of  $P_{g-1}(\lambda)$  does not cross the black open domains (but it can touch their boundaries).*

We associate an admissible polynomial with an admissible divisor. Denote this map by  $\Pi : D \rightarrow P_{g-1}(\lambda)$ . The inverse map  $\Pi^{-1}$  is multivalued. If  $P_{g-1}(\lambda)$  is an admissible polynomial, then the equation (25) has  $2g+1$  roots. One of them is the point  $(0,0)$ , other  $2g$  roots form  $g$  pairs. To define an admissible divisor we have to choose one point from each pair: therefore we have at most  $2^g$  possibilities depending on the number of real points. In the generic case we have no real roots, so the number is equal to  $2^g$  in this case.

It follows immediately from Lemma 3 that the set of all admissible divisors is compact.

We already know that the set of all admissible divisors consists of  $2^m$  connected components. But our knowledge of this fact is noneffective yet: it simply follows from the number of real components found by the different authors many years ago (see the previous section). We are going to describe real components effectively in terms of the admissible polynomials.

Let us associate with each admissible polynomial a **Topological Type**, i.e. a collection of  $m$  numbers  $s_k = \pm 1$ ,  $k = 1, \dots, m$  defined by the following rule:

$$\begin{aligned} s_k &= 1 \text{ if } P_{g-1}(\lambda) \geq f_+(\lambda) \text{ as } E_{2k} \leq \lambda \leq E_{2k-1}, \\ s_k &= -1 \text{ if } P_{g-1}(\lambda) \leq f_-(\lambda) \text{ as } E_{2k} \leq \lambda \leq E_{2k-1}. \end{aligned}$$

**Lemma 4** *Each topological type is presented by the convex subset in the space of polynomials  $P_{g-1}$ .*

Proof. Each topological type is defined by the system of linear inequalities as it follows from the definition above. Lemma is proved.

**Lemma 5** *Each of these  $2^m$  connected components is non-empty and depends continuously on the branching points  $E_1, \dots, E_{2g}$ .*

It is sufficient for the proof to point out, that for any collection of  $m$  numbers  $s_k = \pm 1$ ,  $k = 1, \dots, m$  the polynomial

$$P_{g-1}(\lambda) = \sqrt{-E_1 - E_{2m}}(-1)^{s_1} \prod_{k=2}^m (\sqrt{E_{2k-2}E_{2k-1}} - (-1)^{s_k s_{k-1}} \lambda) \prod_{k=m+1}^g (\sqrt{E_{2k-1}E_{2k}} - \lambda) \quad (28)$$

is admissible and belongs to the appropriate connected component.

The properties of the polynomial (28) can be easily derived from the following elementary inequalities.

Let  $a, b$  be a pair of negative real numbers,  $a < b$ . Then:

$$\begin{aligned} \left| \sqrt{\frac{(x-a)(x-b)}{x}} \right| &> \sqrt{-a-b} \quad \text{as } x > 0, \\ \left| \sqrt{\frac{(x-a)(x-b)}{x}} \right| &< \sqrt{-a-b} \quad \text{as } a < x < b. \end{aligned} \quad (29)$$

Let  $a, b$  be a pair of negative real numbers,  $a < b$ . Then

$$\begin{aligned} \left| \sqrt{(x-a)(x-b)} \right| &> \left| (\sqrt{ab} \pm x) \right| \quad \text{as } x > 0, \\ \left| \sqrt{(x-a)(x-b)} \right| &< \left| (\sqrt{ab} \pm x) \right| \quad \text{as } b < x < 0, \\ \left| \sqrt{(x-a)(x-b)} \right| &< \left| (\sqrt{ab} \pm x) \right| \quad \text{as } x < a. \end{aligned} \quad (30)$$

Let  $a, b$  be a complex number with non-zero imaginary part,  $f(x) = \sqrt{(x-a)(x-\bar{a})}$ . Then

$$\begin{aligned} \left| \sqrt{(x-a)(x-\bar{a})} \right| &> \left| (\sqrt{a\bar{a}} - x) \right| \quad \text{as } x > 0, \\ \left| \sqrt{(x-a)(x-\bar{a})} \right| &< \left| (\sqrt{a\bar{a}} - x) \right| \quad \text{as } x < 0. \end{aligned} \quad (31)$$

Combining the inequalities (29)-(31) we are coming to the conclusion:

For the polynomial  $P_{g-1}(\lambda)$  defined by the formula (28) we have the following inequalities:

$$\begin{cases} \left| \frac{\sqrt{R(\lambda)}}{\lambda} \right| > |P_{g-1}(\lambda)| & \text{as } x > 0, \\ \left| \frac{\sqrt{R(\lambda)}}{\lambda} \right| < |P_{g-1}(\lambda)| & \text{as } x < 0 \text{ and the left-hand side is real.} \end{cases} \quad (32)$$

Therefore the polynomial  $P_{g-1}(\lambda)$  is admissible. Lemma is proved.

The following property of admissible divisors is essential:

**Lemma 6** *Projections of the points of admissible divisors to the  $\lambda$ -plane could not lie in the segments  $[E_1, E_0]$ ,  $[E_3, E_2], \dots, (-\infty, E_{2m}]$ . Moreover, it is possible to choose an open neighborhood of these segments such, that the projections of these points could not lie in it.*

*Proof of Lemma 6.* If  $(\lambda_k, \mu_k)$  is a point of admissible divisor such, that

1.  $\lambda \in \mathbb{R}$ .
2.  $E_{2j} < \lambda_k < E_{2j-1}$ ,  $1 \leq j \leq m$ ,

then  $R(\lambda_k) < 0$  and  $\mu_k$  is a pure imaginary number. But from the description of admissible divisors we have  $\mu_k = \lambda P_{g-1}(\lambda_k)$  and  $\mu_k$  should be real. We have obtained a contradiction.

Assume now, that  $\lambda_k = E_j$  where  $E_j$  is one of real branching points,  $j \neq 0$ . Then  $\mu_k = 0$  and  $P_{g-1}(\lambda_k) = 0$ . In the neighborhood of  $E_j$  the functions  $f_{\pm}(\lambda)$  behaves like  $\sqrt{c(\lambda - E_j)}$ ; therefore the graph of  $P_{g-1}(\lambda)$  intersects the corresponding black domain. We have obtained a contradiction.

The existence of such a neighborhood follows from the compactness of the set of admissible divisor.

Lemma 6 is proved.

**Lemma 7** *Let projection of a point  $\gamma_s = (\lambda_s, \mu_s)$  of an admissible divisor to the  $\lambda$ -plane lie in the segments  $[E_{2k}, E_{2k-1}]$ ; then we have either  $\mu_s < 0$  for the case  $s_k = 1$  or  $\mu_s > 0$  if  $s_k = -1$ .*

*Proof of Lemma 7.* By definition  $s_k$  is the sign of the polynomial  $P_{g-1}(\lambda_s)$  at this interval. Taking into account that  $\mu_s = \lambda_s P_{g-1}(\lambda_s)$  and  $\lambda_s < 0$  we complete the proof.

## 4 Topological charge.

We call the spectral curve  $\Gamma$  **stable** if all branching point are real, i.e.  $m = g$ . It is convenient for the stable  $\Gamma$  to choose a real canonical basis of cycles in the following form.




Fig 2.

Let us draw the following system of cuts  $[E_1, E_0], [E_3, E_2], \dots, (-\infty, E_{2g}]$ . After making these cuts we get 2 sheets. It is specific property of the stable case. Let us denote the sheet containing the line  $\lambda \in \mathbb{R}, \lambda > 0, \mu > 0$  by  $G_+$  and the second sheet by  $G_-$  (see Fig 2).

For the general real case we have exactly  $2m$  negative branching points  $E_1, \dots, E_{2m}$  and  $g - m$  complex adjoint pairs  $E_{2j-1}, E_{2j} = \bar{E}_{2j-1}, j = m + 1, \dots, g$ . The first  $m$   $a$ -cycles should be chosen as in the stable case. The last  $g - m$   $a$ -cycles should be chosen as coverings on  $\Gamma$  over the path on the  $\lambda$ -plane connecting the points  $E_{2j-1}$  and  $E_{2j}$  for  $j > m$ . All these path should not meet each other. All of them should cross positive part of the line  $\lambda > 0$  in one point  $\kappa_j$  (i.e. in two points of Riemann surface) such that  $\kappa_{m+1} < \kappa_{m+2} < \dots < \kappa_g$ . This basis depends on the order of the complex conjugate pairs only (see Fig 3).




Fig 3.

For the complex conjugate pairs of branching points  $E_{2j}, E_{2j-1}, j > m$ , we make the cuts connecting each of these points with  $i\infty$  or  $-i\infty$  such that the cuts do not intersect the real line. As before we denote the sheet containing the line  $\lambda \in \mathbb{R}, \lambda > 0$   $\mu > 0$  by  $G_+$  and the second sheet by  $G_-$ .

Denote by  $\omega^1, \dots, \omega^{g-1}$  the canonical basis of holomorphic differentials in  $\Gamma$ ,

$$\omega^k = i \frac{\sum_{j=0}^{g-1} D_j^k \lambda^j}{\mu} d\lambda, \quad D_j^k \in \mathbb{R} \quad (33)$$

**Theorem 1** *For any real solution the density of topological charge along the variable  $x$  is given by the formula*

$$\bar{n} = \frac{1}{2} \sum_{k=1}^m (-1)^{k-1} s_k \left( D_{g-1}^k + \frac{D_0^k}{\sqrt{\prod_{j=1}^{2g} E_j}} \right). \quad (34)$$

**Proof.** It follows from the  $\Theta$ -functional formulas that the function  $\exp\{iu(x, t)\}$  is a regular function on a real  $g$  dimensional torus restricted to a two-



dimensional linear subspace. Each such torus corresponds to the fixed topological type; it is a real component of the Jacoby torus  $T^{2g} = J(\Gamma)$  with respect to an antiholomorphic involution. Therefore we can apply Lemma 1.

Our calculation consists of two steps. At the first step we calculate the coefficients of the vector  $\vec{v}$ , and at the second step we calculate the topological charges along the basic cycles.

The first step is rather standard (see, for example, [8]). Dubrovin equations (16), (18) can be written in the following form

$$\frac{\partial \lambda_k(\xi, \eta)}{\partial \xi} = -\text{res}_{\gamma=\gamma_k(\xi, \eta)}(\partial_\xi \log \psi_1(\gamma, \xi, \eta))d\lambda, \quad (35)$$

$$\frac{\partial \lambda_k(\xi, \eta)}{\partial \eta} = -\text{res}_{\gamma=\gamma_k(\xi, \eta)}(\partial_\eta \log \psi_1(\gamma, \xi, \eta))d\lambda, \quad (36)$$

From (35), (36) we obtain following formulas for the  $\xi, \eta$  derivatives of the Abel transform of the divisor:

$$\frac{\partial A^k(\vec{\gamma})}{\partial \xi} = -\sum_k \text{res}_{\gamma=\gamma_k(\xi, \eta)} \omega^k \partial_\xi \log \psi_1(\gamma, \xi, \eta) = \text{res}_{\gamma=\infty} \omega^k \partial_\xi \log \psi_1(\gamma, \xi, \eta), \quad (37)$$

$$\frac{\partial A^k(\vec{\gamma})}{\partial \eta} = -\sum_k \text{res}_{\gamma=\gamma_k(\xi, \eta)} \omega^k \partial_\eta \log \psi_1(\gamma, \xi, \eta) = \text{res}_{\gamma=0} \omega^k \partial_\eta \log \psi_1(\gamma, \xi, \eta). \quad (38)$$

Calculating residues in (37), (38) we obtain:

$$\frac{\partial A^k(\vec{\gamma})}{\partial \xi} = 2D_{g-1}^k, \quad (39)$$

$$\frac{\partial A^k(\vec{\gamma})}{\partial \eta} = 2 \frac{D_0^k}{\sqrt{\prod_{j=1}^{2g} E_j}}, \quad (40)$$

and

$$\frac{\partial A^k(\vec{\gamma})}{\partial x} = \frac{1}{2} \left( D_{g-1}^k + \frac{D_0^k}{\sqrt{\prod_{j=1}^{2g} E_j}} \right) = -\frac{1}{2\pi} \oint_{b_k} dp, \quad (41)$$

where  $dp$  denotes a meromorphic differential with zero  $a$ -periods and two second order poles at  $0, \infty$  such, that

$$dp = \begin{cases} \left( \frac{1}{8\sqrt{\lambda}} + o\left(\frac{1}{\lambda}\right) \right) d\lambda & \text{as } \lambda \rightarrow \infty, \\ \left( \frac{1}{8\lambda\sqrt{\lambda}} + o\left(\frac{1}{\lambda}\right) \right) d\lambda & \text{as } \lambda \rightarrow 0 \end{cases} \quad (42)$$

Therefore the density of topological charge is equal to the following expression

$$\bar{n} = \frac{1}{2} \sum_{k=0}^{g-1} \left( D_{g-1}^k + \frac{D_0^k}{\sqrt{\prod_{j=1}^{2g} E_j}} \right) n_k, \quad (43)$$

where  $n_k$  denote the topological charge of the image of the  $k$ -th basic cycle for the given topological component. By definition  $n_k$  is an integer.

To calculate  $n_k$  for a given topological type consider a basic cycle  $\mathcal{A}_k$  on the real component of Jacoby torus. represented by the closed curve  $B_k$  on this torus. The image of this cycle in the surface  $\Gamma$  (i.e. image of it under the inverse Abel map) is a closed oriented curve  $C_k$  (may be consisting from several connected components). This curve is presented geometrically as a union of the divisor points  $D(t_k) = \sum \gamma_j(t_k)$  for all values of the corresponding "time"  $t_k$  along this basic cycle. It is homological to the cycle  $a_k \in H_1(\Gamma, \mathbb{Z})$ . The projection of the last curve  $C_k$  on the  $\lambda$ -plane we denote  $\tilde{C}_k$ . This curve does not touch the closed segments on the real line  $[-\infty, E_{2m}], \dots, [E_3, E_2], [E_1, 0]$ .

**Lemma 8** *The topological charge  $n_k$  along the cycle  $\mathcal{A}_k$  is given by the following formula:*

$$n_k = \tilde{C}_k \circ \mathbb{R}_-, \quad (44)$$

where  $\tilde{C}_k$  is the projection of  $C_k$  to the  $\lambda$ -plane,  $\mathbb{R}_-$  is the negative part of the real line with the standard orientation, and  $\circ$  denotes the intersection index of curves on the  $\lambda$ -plane.

This Lemma follows directly from formula (15).

Assume now, that the cycles  $b_1, b_2, \dots, b_m$  are the ovals lying over the segments  $[E_2, E_1], [E_4, E_3], [E_{2m}, E_{2m-1}]$  respectively. It is easy to check,

that the function  $\mu$  at the sheet  $G_+$  is negative at the intervals  $[E_{4k-2}, E_{4k-3}]$  and positive at the intervals  $[E_{4k}, E_{4k-1}]$ .




Fig 4.

It follows from Lemma 6, that the projection-curve  $\tilde{C}_k$  can cross the half-line  $\mathbb{R}_-$  only strictly inside the intervals  $[E_2, E_1]$ ,  $[E_4, E_3]$ ,  $[E_{2m}, E_{2m-1}]$ . Therefore any time when the curve  $\tilde{C}_k$  crosses the half-line  $\mathbb{R}_-$ , the cycle  $C_k$  on the Riemann surface  $\Gamma$  crosses one of the cycles  $b_1, b_2, \dots, b_m$ . The orientation of the cycles  $b_l$ ,  $l = 1, \dots, m$  coincides with the orientation of  $\mathbb{R}_-$  at the sheet  $G_+$ ; they are opposite at the sheet  $G_-$ . It follows from Lemma 7, that the cycle  $C_k$  always crosses  $b_l$  in the sheet  $G_+$  if  $(-1)^{l-1}s_l > 0$ ; it crosses  $b_l$  in the sheet  $G_-$  if  $(-1)^{l-1}s_l < 0$ . Therefore we proved

**Lemma 9** *The topological charge along the cycle  $\mathcal{A}_k$  on the real torus in the Jacobian variety is equal to the intersection index of the corresponding curve  $C_k$  with the **Counting Cycle of the real component** on the Riemann surface  $\Gamma$*

$$n_k = C_k \circ \left( \sum_{l=1}^m (-1)^{l-1} s_l b_l \right) \quad (45)$$

Let us mention that the cycle  $C_k$  is homotopic to  $a_k$  on  $\Gamma$ . Therefore  $C_k \circ b_l = \delta_{kl}$  and

$$\begin{cases} n_k = (-1)^{k-1} s_k & \text{for } k \leq m \\ n_k = 0 & \text{for } k > m. \end{cases} \quad (46)$$

Combining (43) with (46), we obtain our main formula (34).

The Theorem is proved.

Let us present here a second method for calculating the topological charges along the basic cycles. By Lemma 5 the topological components depend continuously on the branching points. Assuming that the admissible polynomial is given by (32) and all points of the divisor  $D$  lie in the upper half-plane we obtain a family of solutions continuously depending on the branching points. The motion of an individual zero point  $\gamma_j$  with any  $2\pi$ -cyclic time  $\tilde{t}_s$  corresponding to the basic cycle  $\mathcal{A}_s$  on the real torus does not depend continuously on the initial data, because the divisor points may collide with each other; the Dubrovin equations became singular in the collision point. The set of singular divisors has real codimension 2. But we deduce from the  $\Theta$ -functional formulas that all symmetric functions of the divisor points pass through this singularity continuously. All real solutions are nonsingular. Therefore the topological charges along the basic cycles are integers continuously depending on the branching points. We conclude that they are constant.

For the proof of (46) it is sufficient to calculate the number  $n_k$  for one spectral curve  $\Gamma_k$ . Let  $k \leq m$ . Consider the following configuration of the branching points for this curve  $\Gamma_k$ :

$$\begin{cases} E_{2k} = -k - \epsilon_1 & E_{2k-1} = -k + \epsilon_1 \\ E_{2j} = -j - \epsilon_2 & E_{2j-1} = -j + \epsilon_2 & \text{as } j \leq m, j \neq k \\ E_{2j} = j - i\epsilon_2 & E_{2j-1} = j + i\epsilon_2 & \text{as } j > m, \end{cases} \quad (47)$$

where  $\epsilon_2 \ll \epsilon_1 \ll 1$ .

The cycle  $\mathcal{A}_k$  can be realized by the following 1-parametric family of polynomials  $P_{g-1}(\lambda, \beta)$ :

$$P_{g-1}(\lambda, \beta) = (-1)^{k+g-m-1} s_k \beta \prod_{j=1}^{g-1} (\lambda - r_j), \quad (48)$$

where

$$\begin{aligned} r_j &= -j - (-1)^{k+j} s_j s_k 2|j - k| \frac{\epsilon_2}{\epsilon_1} \sqrt{\frac{k}{j}} & \text{as } j < k \\ r_j &= -j - 1 - (-1)^{k+j} s_j s_k 2|j + 1 - k| \frac{\epsilon_2}{\epsilon_1} \sqrt{\frac{k}{j+1}} & \text{as } k \leq j < m \\ r_j &= j + 1 & \text{as } m \leq j < g, \end{aligned} \quad (49)$$

and the parameter  $\beta$  runs through the interval  $[\frac{\epsilon_1}{\sqrt{k}}(1 + o(1)), 2\sqrt{k}(1 + o(1))]$ .

Elementary estimates show, that system (25) has a pair of roots located in a neighborhood of the circle  $|\lambda| = k$  and all other roots form complex conjugate pairs located in small neighborhoods of the point  $r_n$ . The distinguished pair of roots lie in the sheet  $G_+$  if  $(-1)^{k-1} s_k > 0$  and in  $G_-$  if  $(-1)^{k-1} s_k < 0$ . Therefore we proved (46) for  $k \leq m$ .

To calculate  $n_l$  with  $l > m$  it is sufficient to consider the same spectral curve as above. Let us assume  $\beta = 1$  and consider a small variation of the point  $r_{l-1}$ . It gives us the cycle  $a_l$ . Variation of the function  $u(\xi, \eta)$  along this cycle is very small, therefore  $n_l = 0$ .

The second proof of the Theorem is completed.

## 5 Appendix 1. The Cherednik differential for the generic periodic real SG solutions.

Let us explain here why the sufficient reality conditions from the Section 2.3 are in fact necessary for the generic smooth  $x$ -periodic solutions. Let us remind, that in the SG theory we call a potential  $u(x, t)$  periodic if the function  $\exp\{u(x, t)\}$  is periodic. Therefore the operator

$$L(\lambda) = \partial_x - 1/4(U + V)$$

has a Bloch solution

$$L(\lambda)\Psi = 0, \quad \hat{T}\Psi_{\pm}(\lambda, x, t) = \Psi_{\pm}(\lambda, x + T, t) = \exp\{\pm ip(\lambda)T\}\Psi_{\pm}(\lambda, x, t)$$

where the differential  $dp$  on the Riemann surface has been defined above in the finite-gap case. This formula works for all complex values of  $\lambda$  except branching points where the shift operator  $\hat{T}$  has the Jordan cells. The Bloch vector-function  $\Psi_{\pm}(\lambda, x)$  is in fact a meromorphic one-valued vector-function  $\Psi(\gamma, x)$  on the Riemann surface  $\Gamma$  whose points are exactly the pairs  $\gamma =$

$(\lambda, \pm) \in \Gamma$  or  $\gamma = (\lambda, \mu)$  for  $\mu = \exp\{ipT\}$ . This surface is obviously 2-sheeted, i.e. hyperelliptic. For the real solutions this surface has a pair of commuting involutions (holomorphic and antiholomorphic):

$$\begin{aligned}\sigma(\lambda, +) &= (\lambda, -), \quad \text{or} \quad \sigma(\lambda, \mu) = (\lambda, \mu^{-1}), \\ \tau(\lambda, \mu) &= (\bar{\lambda}, \overline{\mu^{-1}}).\end{aligned}$$

The action of  $\sigma\tau$  on the Bloch solutions is given by the following explicit formula

$$\begin{pmatrix} \psi_1(\sigma\tau\gamma, x, t) \\ \psi_2(\sigma\tau\gamma, x, t) \end{pmatrix} = \frac{1}{\psi_2(\gamma, 0, 0)} \begin{pmatrix} \overline{\psi_2(\gamma, x, t)} \\ -\lambda\psi_1(\gamma, x, t) \end{pmatrix} \quad (50)$$

**Statement.** The Riemann surface  $\Gamma$  always has the points  $0, \infty$  as its branching points. For the real smooth  $x$ -periodic SG solutions  $u(x, t)$  this surface has no real positive branching points. Let this surface be nonsingular. Then we define a "Cherednik differential"  $\Omega$ : by definition, its zeroes should coincide with the full set of poles of the Bloch function with the proper normalization

$$\psi_{1,\pm}(\lambda, x, t) = 1 \quad \text{for} \quad x = 0, t = 0, \quad (51)$$

and their  $\tau$ -conjugate points. It has also simple poles in the points  $0, \infty$ . This differential can be written in the following form:

$$\Omega = \frac{\psi_1(\gamma, 0, 0)\psi_2(\sigma\gamma, 0, 0)}{\psi_1(\gamma, 0, 0)\psi_2(\sigma\gamma, 0, 0) - \psi_1(\sigma\gamma, 0, 0)\psi_2(\gamma, 0, 0)} d\lambda. \quad (52)$$

Let us check that the differential  $\Omega$  defined by (52) is holomorphic outside the points  $0, \infty$  and has the correct set of zeroes. We see, that the form  $\Omega$  does not depend on the normalization of the Bloch function  $\Psi$ . Therefore we may assume, that any locally holomorphic normalization is used. Then the denominator of formula (52) has zeroes exactly at the points where the solutions  $\Psi(\gamma, x, t)$  and  $\Psi(\sigma\gamma, x, t)$  are linearly dependent, i.e. at the branching points. The surface  $\Gamma$  is nonsingular, therefore these zeroes are simple and the form  $\Omega$  has no singularities at the branching points.

The zeroes of the form  $\Omega$  are the zeroes of the first component of the Bloch function  $\psi_1(\gamma, 0, 0)$  plus the zeroes of the second multiplier  $\psi_2(\sigma\gamma, 0, 0)$ . It follows from formula (50) that zeroes of the function  $\psi_2(\sigma\gamma, 0, 0)$  are  $\tau$ -conjugate to the zeroes of the function  $\psi_1(\gamma, 0, 0)$ . Taking into account that the poles of the Bloch function normalized by the condition (51) are exactly the zeroes of  $\psi_1(\gamma, 0, 0)$ , we complete the proof.

Let  $\lambda$  be a real positive number. Then the restriction of the shift operator  $\hat{T}$  to the zero eigenspace of the operator  $L(\lambda)$  is unitary. Therefore  $\hat{T}$  and has no Jordan cells, and  $\lambda$  is not a branching point.

## 6 Appendix 2. Numerical simulations.

During the work on this problem the following numerical simulations were made:

1. The topological charges along the basic cycles on the real part of the Jacoby torus were calculated for all topological types.
2. The densities of topological charge for the  $x$ -dynamics and for the  $\xi$ -dynamics were directly calculated.

For numerical calculation of topological charges  $n_k$  along the basic cycles we used the following approach. Assume, that the Baker-Akhiezer function  $\Psi(\gamma, \xi, \eta)$  has  $2g$  additional essential singularities at the points  $(\lambda, \mu) = (\beta_s, \pm\sqrt{R(\beta_s)})$  with the following asymptotics:

$$\Psi(\gamma, \xi, \eta) = O(1)e^{-i\frac{\pm\sqrt{R(\beta_s)}\tilde{t}_s}{\lambda - \beta_s}} \quad (53)$$

where  $\beta_s$ ,  $s = 1, 2, \dots, g$  are some real numbers. It is rather easy to check, that these flows commute with the SG-dynamics. They correspond to the motion along straight lines in the real components of the Jacoby torus:

$$\frac{\partial A^k(\vec{\gamma})}{\partial \tilde{t}_s} = W_s^k = 2 \sum_{l=0}^{g-1} D_l^k \beta_s^l. \quad (54)$$

These flows are non-local in terms of the Sine-Gordon potential  $u$ , but the corresponding Dubrovin equations are well-defined and have rational right-

hand sides:

$$\left\{ \begin{array}{l} \frac{\partial \lambda_l}{\partial \tilde{t}_s} = -2i \frac{\mu_l \prod_{j \neq l} (\beta_s - \lambda_j)}{\prod_{j \neq l} (\lambda_l - \lambda_j)} \\ \frac{\partial \mu_l}{\partial \tilde{t}_s} = -i \frac{R'(\lambda_l) \prod_{j \neq l} (\beta_s - \lambda_j)}{\prod_{j \neq l} (\lambda_l - \lambda_j)} \\ \frac{\partial u}{\partial \tilde{t}_s} = -2 \sum_l \frac{\mu_l \prod_{j \neq l} (\beta_s - \lambda_j)}{\lambda_l \prod_{j \neq l} (\lambda_l - \lambda_j)}. \end{array} \right. \quad (55)$$

To avoid singularities in the flows (55) generated by collisions of divisor projections  $\lambda_k(\xi, \eta, \tilde{t}_s)$  with the points  $\beta_s$  we have chosen the points  $\beta_s$  in the forbidden intervals:

$$\beta_s = \frac{1}{2}(E_{2s-1} + E_{2s-2}), \quad s = 1, \dots, g. \quad (56)$$

It follows from equation (54), that the linear combination of non-local flows

$$\frac{\partial}{\partial \nu_k} = \sum_{l=1}^g (W^{-1})_k^l \frac{\partial}{\partial \tilde{t}_l} \quad (57)$$

where  $W^{-1}$  denote the inversion of matrix  $W$ , corresponds to the motion along the basic cycle  $\mathcal{A}_k$  in the real torus. Integrating (57) numerically we obtained the topological charges along the basic cycles.

## 7 Appendix 3. Fourier Transform on the Riemann Surfaces: The continuous analog of the Krichever-Novikov bases.

Let us invent here a continuous analog of the so-called “Krichever-Novikov bases” on the Riemann surfaces defined and used by the authors for the needs of the quantum string theory in the late 80s—see the first work in this



series [27]. These bases have been considered as a natural analog of the Fourier-Laurent bases in complex analysis.

Let us consider the following set of data: a nonsingular Riemann surface  $\Gamma$  of the genus  $g$  with marked point  $\infty \in \Gamma$  and selected local parameter near this point  $z = k^{-1}$ ,  $z(\infty) = 0$ . We construct a function  $\psi^0 = \psi(P, x)$  holomorphic on  $\Gamma \setminus \infty$  and exponential near the infinite point:

$$\psi(z, x) = k^g \exp\{kx\} \left(1 + \sum_{i>0} \eta_i(x) k^{-i}\right), \quad \eta_0 = 1$$

It is obviously a partial case of the general Krichever KP-construction for the case where the divisor of poles is equal to  $g\infty = D$ . This case leads to the singular solutions of the KP-type systems. Therefore it is out of use in the physical soliton theory. However, the examples of this kind appeared in the works of XIX century as the so-called Lamé operator. It appears as a result of the separation of variables in the Laplace-Beltrami operator on the 3-axis ellipsoid. The spectral theory on the closed segment between two singularities has been studied by the classical people like Hermit and others:

**Example.** For the genus  $g = 1$  we have:

$$\psi(z, x) = \frac{\sigma(z-x)}{\sigma(z)\sigma(x)} \exp\{x\zeta(z)\}$$

This function satisfies to the Lamé equation

$$(\partial_x^2 - 2\wp(x))\psi = \wp(z)\psi(z, x)$$

We construct also a differential form  $\psi^1 = \phi(P, x)dz$  on  $\Gamma$  such that it is holomorphic outside of the infinite point and has following asymptotics near it:

$$\phi(P, x)dz = z^{g-2} \exp\{-kx\}(1 + o(1))dz.$$

Consider a  $x$ -independent second kind differential  $dp$  on the surface  $\Gamma$  with the unique pole of the second order in the point  $\infty \in \Gamma$  and with asymptotics:

$$dp = dk(1 + o(1)), \quad k = z^{-1} \rightarrow \infty$$

We require that this differential has following property for all 1-dimensional cycles on the Riemann surface:

$$\operatorname{Re} \oint_A dp = 0, \quad A \in H_1(\Gamma, \mathbb{Z})$$

Therefore the function  $\operatorname{Re} p = \tau$  is an one-valued real harmonic function  $\tau(P)$  on the Riemann surface with the unique pole in the point  $\infty$ . This function will be called “time” by the analogy with the string theory [27]. All its levels enter the point  $\infty$  twice.

**Example.** For the genus equal to 1  $g = 1$  and real algebraic curves we have the periods  $2\omega \in R$  and  $2i\omega' \in iR$ . The function defined above has a form:

$$p(z) = \zeta(z) - \frac{\eta}{\omega}z, \quad dp(z) = \left(-\wp(z) - \frac{\eta}{\omega}\right)dz$$

where  $\eta = \zeta(\omega)$  (see the book [28]).

**Problem.** How to construct Harmonic Analysis on the levels  $\tau = \text{const}$  and in the complex domains  $c_1 < \tau < c_2$  using the special Baker-Akhiezer function  $\psi(P, x)$  described above? Which multiplicative properties have the basic functions  $\psi(P, x) = \psi_x(P)$  depending on  $x$  as parameter?

**Proposition.** The Fourier-Laurent decomposition formula for the function  $f(P)$  has a form:

$$f(P) = \int \left\{ 1/2\pi \oint_{\tau=\text{const}} \phi(P', x) f(P') dz \right\} \psi(P, x) dx$$

here the points  $P, P'$  belong to the level  $\tau = \text{const}$  or to the domain  $c_1 < \tau < c_2$ . All connectivity components should enter this integral. This formula valid for the class of functions with the proper decay near  $\infty$ . We shall publish more precise statement later.

We consider here the subscript  $x$  as a continuous analog of the discrete  $n \in Z$  from the work [27]. Therefore we expect to have here some natural analog of the “almost grading property” discovered in [27]:

**Theorem 2 .** *Let  $x, y \neq 0$ . There exists a differential operator  $L$  in the variable  $x$  of the order  $g$  with coefficients dependent on the both variables  $x, y$  such that the following **Almost Graded Commutative Associative Ring Structure** is defined by the formula*

$$\psi(P, x)\psi(P, y) = L\psi(P, x + y)$$

$$L = \partial_x^g + [\eta(x) + \eta(y) - \eta(x + y)]\partial_x^{g-1} + \dots$$

*The coefficients of this operator have poles for the points  $x = 0$  or  $y = 0$ .*

In the example above where  $g = 1$  we have

$$L = \partial_x - (\zeta(x) + \zeta(y) - \zeta(x + y))$$

In the next work we shall present more detailed description of this idea including the higher tensor generalization  $\psi^k$ , the decomposition theorems for the functions and tensors on the contours  $\tau = \text{const}$  and domains  $c_1 < \tau < c_2$  following the program of the work [27].

## References

- [1] Novikov S.P. The periodic problem for the Korteweg-de Vries equation. I// Funkts. Anal. i ego Pril., 1974, V. 8, No. 3, P. 54–66.; English translation: Funct. Anal. Appl., 1974, V. 8, P. 236–246.
- [2] Dubrovin B.A., Novikov S.P. Periodic and conditionally periodic analogs of the many-soliton solutions of the Korteweg-de Vries equation// Zh. Eksper. Teoret. Fiz., 1974, V. 67, No. 6, P. 2131–2144; English translation: Sov. Phys-JETP, 1974, V. 40, P. 1058–1063.
- [3] Dubrovin B.A., Novikov S.P. A periodicity problem for the Korteweg-de Vries and Sturm-Liouville equations. Their connections with algebraic geometry// Dokl. Akad. Nauk SSSR, 1974, V. 219, No. 3, P. 531–534; English translation: Sov. Math. Dokl., 1974, V. 15, P. 1597–1601.
- [4] Dubrovin B.A. Periodic problem for the Korteweg-de Vries equation in the class of finite-band potentials// Funkts. Anal. i ego Pril., 1975, V. 9, No. 3, P. 41–51. English translation: Funct. Anal. Appl., 1975, V. 9, P. 215–223.
- [5] Its A.R., Matveev V.B. Hill operators with finitely many gaps// Funkts. Anal. i ego Pril., 1975, V. 9, No. 1, P. 69–70; English translation: Funct. Anal. Appl., 1975, V. 9, P. 65–66.
- [6] Lax P.D. Periodic solutions of the KdV equation// Comm. Pure Appl. Math. 1975, V. 28, P. 141–188.
- [7] McKean H.P., van Moerbeke P. The spectrum of Hill’s equation// Invent. Math., 1975, V. 30, No. 3, P. 217–274.

- [8] Dubrovin B.A., Matveev V.B., Novikov S.P. Non-linear equations of Korteweg-de Vries type, finite-zone linear operators and Abelian varieties// *Uspekhi Mat. Nauk*, 1976, V. 31, No. 1, P. 55–136; English translation: *Russ. Math. Surveys*, 1976, V. 31, No 1, P. 59–146.
- [9] Flaschka H., McLaughlin D.W. Canonically conjugate variables for the Korteweg-de Vries equation and the Toda lattice with periodic boundary conditions// *Progr. Theoret. Phys.* 1976, V. 55, No. 2, P. 438–456.
- [10] Novikov S.P., Veselov A.P. On Poisson brackets compatible with algebraic geometry and Korteweg-de Vries dynamics on the set of finite-zone potentials// *Dokl. Akad. Nauk SSSR*, 1982, V. 266, No. 3, P. 533–537; English translation: *Sov. Math. Dokl.*, 1982, V. 26, P. 357–362.
- [11] Krichever I.M. An algebraic-geometric construction of the Zakharov-Shabat equations and their periodic solutions// *Dokl. Akad. Nauk SSSR*, 1976, V. 227, No. 2, P. 291–294; English translation: *Sov. Math. Dokl.*, 1976, V. 17, P. 394–397.
- [12] Dubrovin B.A., Krichever I.M., Novikov S.P. The Schrödinger equation in a periodic field and Riemann surfaces// *Dokl. Akad. Nauk SSSR*, 1976, V. 229, No. 1, P. 15–18; English translation: *Sov. Math. Dokl.*, 1976, V. 17, P. 947–951.
- [13] Novikov S.P., Veselov A.P. Finite-zone, two-dimensional Schrödinger operators. Potential operators// *Dokl. Akad. Nauk SSSR*, 1984, V. 279, No. 4, P. 784–788. English translation: *Sov. Math. Dokl.*, 1984, V. 30, P. 705–708; Finite-zone, two-dimensional potential Schrödinger operators. Explicit formulas and evolution equations// *Dokl. Akad. Nauk SSSR*, 1984, V. 279 No. 1, P. 20–24; English translation: *Sov. Math. Dokl.*, 1984, V. 30, P. 588–591.
- [14] Krichever I.M., Novikov S.P. Holomorphic vector bundles over Riemann surfaces and the Kadomtsev-Petviashvili equation. I// *Funkts. Anal. i ego Pril.*, English translation: *Funct. Anal. Appl.*, 1978, V. 12, P. 276–286.
- [15] Dubrovin B.A., Natanzon S.M. Real two-zone solutions of the sine-Gordon equation// *Funkts. Anal. i ego Pril.*, 1982, V. 16, No. 1, P. 27–43; English translation: *Funct. Anal. Appl.*, 1982, V. 16, P. 21–33.

- [16] Ercolani N.M., Forest M.G. The geometry of real Sine-Gordon wave-trains// Comm. Math. Phys. 1985. V. 99, No. 1. P. 1–49.
- [17] Dubrovin B.A., Novikov S.P. Algebro-geometrical Poisson brackets for real finite-zone solutions of the Sine-Gordon equation and the nonlinear Schrödinger equation// Dokl. Akad. Nauk SSSR, 1982, V. 267, No. 6, P. 1295–1300; English translation: Sov. Math. Dokl., 1989, V. 26, No. 3, P. 760–765.
- [18] Novikov S.P. Algebrotopological approach to the reality problems. Real action variables in the theory of finite-zone solutions of the Sine-Gordon equation// Zap. Nauchn. Sem. LOMI: Differential geometry, Lie groups and mechanics, VI., 1984, V. 133. P. 177–196; English translation: J. Soviet Math., 1985, V. 31, No. 6, P. 3373–3387.
- [19] Grinevich P.G., Novikov S.P. Real finite-gap Sine-Gordon solutions: a formula for topological charge (in Russian)// Uspekhi. Mat. Nauk, 2001, V. 56, No. 5, P. 181–182.; English translation: to appear in Russ. Math. Surveys, V. 56, No. 5.
- [20] Lamb G.L. Analytical description of ultrashort optical pulse propagation in a resonant medium// Rev. Mod. Phys, 1971, V. 43, P. 99–124.
- [21] Ablowitz M.J., Kaup D.J., Newell A.S., Segur H. Method for solving the Sine-Gordon equation// Phys. Rev. Lett., 1973. V. 30. No. 25. P. 1262–1264.
- [22] Novikov S.P., Veselov A.P. Exactly solvable two-dimensional Schrödinger operators and Laplace transformations. In: Solitons, geometry, and topology: on the crossroad// Amer. Math. Soc. Transl. Ser. 2, 1997, V. 179, P. 109–132.
- [23] Kozel V.A., Kotlyarov V.P. Almost periodic solutions of the equation  $u_{tt} - u_{xx} + \sin u = 0$ . (Russian)// Dokl. Akad. Nauk Ukrain. SSR Ser. A, 1976, No. 10, P. 878–881.
- [24] Its A.R., Kotlyarov V.P. Explicit formulas for solutions of the Nonlinear Schrödinger equation. (Russian)// Dokl. Akad. Nauk Ukrain. SSR, Ser. A, 1976, No. 11, P. 965–968;

- [25] Cherednik I.V. Reality conditions in “finite-zone” integration// Dokl. Akad. Nauk SSSR, 1980, V. 252, No. 5, P. 1104–1108; English translation: Sov. Phys. Dokl., 1980, V. 25, P. 450–452.
- [26] Natanzon S.M. Moduli of real algebraic surfaces, and their superanalogues. Differentials, spinors and Jacobians of real curves// Uspekhi Mat. Nauk, 1999, V. 54, No. 6, P. 3–60; English translation: Russ. Math. Surveys, 1999, V. 54, No. 6, P. 1091–1147.
- [27] Krichever I.M., Novikov S.P. Algebras of Virasoro type, Riemann surfaces and the structures of soliton theory// Funkts. Anal. i ego Pril., 1987, V. 21, No. 2, P. 46–63; English translation: Funct. Anal. Appl., 1987, V. 21, No 2.
- [28] Higher transcendental functions. Volume 3. Based in part on notes left by Harry Bateman and completed by the staff of the Bateman manuscript project. Director Arthur Erdélyi. New-York, Toronto, London, Mc Graw-Hill book company, Inc. 1955.